

Complex Retained Ballistic Foreign Bodies of the Nasopharynx and Ethmoid Sinus: A Two-Case Series and Review

Nongthombam Bidyananda Singh¹, Sauradeep Das², Manu CB³, Suvamoy Chakraborty⁴

Received on: 3rd August 2025; Accepted on: 17th August; Published on: 5th September 2025

ABSTRACT

Background: Retained ballistic foreign bodies in anatomically intricate regions such as the nasopharynx or paranasal sinuses are exceptionally rare and pose diagnostic and therapeutic challenges.

Cases: We present two cases in 49-year-old males: Case 1 involved a bullet lodged in the posterior nasopharyngeal wall with multiple tooth fragments embedded in the tongue following intraoral gunshot trauma; Case 2 featured a metallic fragment retained in the left posterior ethmoid sinus after a self-inflicted nasal gunshot wound.

Management and Outcome: Both patients underwent emergency surgical intervention—open retrieval and tracheostomy for Case 1, endoscopic sinus surgery with septoplasty for Case 2—and recovered uneventfully.

Conclusion: These cases underscore the critical role of high resolution computed tomography, multidisciplinary planning, and tailored surgical approaches in successfully managing rare upper aerodigestive ballistic injuries.

Keywords: Ballistic trauma; retained foreign body; nasopharynx; ethmoid sinus; endoscopic sinus surgery; Computed tomography scan imaging; gunshot wound

INTRODUCTION

Firearm injuries to the head and neck constitute approximately 6% of all gunshot wounds and frequently result in complex trauma due to the high density of vital structures.⁶ Retained projectiles or fragments in confined spaces—such as the nasopharynx and paranasal sinuses—are exceedingly uncommon, often following unpredictable ballistic trajectories after deflection by bone or teeth.^{1,4,7,11} These retained foreign bodies may remain asymptomatic or present with nonspecific features, delaying diagnosis and treatment. Potential complications include chronic inflammation, granuloma, infection, and systemic lead toxicity.^{2,3,5,7,12} High resolution computed tomography scan (CT) is indispensable for localizing radiopaque material and guiding surgical planning.⁵ Herein, we describe two rare cases of retained ballistic foreign bodies—one in the nasopharynx with lingual dental fragments, and one in the posterior ethmoid sinus—highlighting their presentation, management, and outcomes.

Junior Resident¹, Senior Resident², Assistant Professor³, Professor and HOD⁴

Department of ENT¹⁻⁴

North Eastern Indira Gandhi Regional Institute of Health & Medical Sciences, Shillong¹⁻⁴

Corresponding Author: Nongthombam Bidyananda Singh
Junior Resident, Department of ENT

North Eastern Indira Gandhi Regional Institute of Health & Medical Sciences, Shillong

Email ID: nongaboy5@gmail.com

Source of funding: None

Conflict of interest: None declared

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: sabarish@krishers.com

How to cite this article: Nongthombam B, Sauradeep D, Manu CB, Suvamoy C. Complex Retained Ballistic Foreign Bodies of the Nasopharynx and Ethmoid Sinus: A Two-Case Series and Review. *J Otolaryngol Head Neck Surg.* 2025; 1(2):1-24

Case 1: Nasopharyngeal Bullet with Lingual Tooth Fragments

A 49-year-old man presented three days after a self-inflicted intraoral gunshot wound, complaining of severe oropharyngeal pain, drooling, and inability to swallow. Examination revealed missing mandibular premolars and lateral tongue lacerations. CT scan showed a bullet lodged in

Access this article online	
Website: https://ajohns.net/	Quick Response Code 
DOI: 10.66243/Krishers-AJOHNS-014	

the posterior nasopharyngeal wall and multiple hyperdense fragments within tongue musculature (Figure 1 & 2).



Figure 1: (A) X-ray antero-posterior and (B) lateral view of head and neck showing foreign body (bullet) in nasopharynx (red arrow)




Figure 2: (A) and (B) Computed tomography scan axial view showing foreign body (bullet) in nasopharynx (red arrow)




Figure 3: (A) Intra-operative picture showing tongue after exploration and (B) intra-operative X-ray showing residual foreign body in tongue

Under general anesthesia, a diagnostic nasal endoscopy was done. The bullet was difficult to locate intraoperatively, using a transoral approach, a blunt finger dissection was done to locate the foreign body in the posterior wall of the nasopharynx and was brought to the right side, wherein a dissection was done to retrieve the bullet via the nasal cavity under endoscopic visualization. Figure 1C. Four dental fragments were removed from the tongue. Intraoperative radiography confirmed a

minute residual fragment in the tongue, which was deemed irretrievable. A tracheostomy secured the airway in view of tongue edema by the end of the procedure (Figure 3). The patient's postoperative course was uneventful; decannulation occurred on postoperative day 7, with full symptomatic resolution uneventfully.

Case 2: Posterior Ethmoid Sinus Retained Fragment

Another 49-year-old man presented three days after a self-inflicted nasal gunshot injury, reporting nasal pain and epistaxis. Physical examination revealed a healed nasal dorsum laceration and septal deviation. CT demonstrated a displaced nasal bone fracture and a metallic fragment within the left posterior ethmoid sinus, with an additional subcutaneous piece adjacent to the nasal bone (Figure 4).




Figure 4: Computed Tomography of nose and paranasal sinus region showing foreign body in left posterior ethmoidal sinus in axial view (A), coronal view (B) and sagittal view (C)

Diagnostic nasal endoscopy confirmed septal deviation and blood clot on left side. The patient underwent functional endoscopic sinus surgery (FESS) combined with septoplasty; the ethmoidal fragment was extracted without complications. Follow-up endoscopy showed a well healed sinonasal cavity with no retained material.

DISCUSSION

Retained ballistic foreign bodies in the nasopharynx or paranasal sinuses account for <1% of head and neck gunshot wounds.¹¹ Case 1 illustrates how intraoral trajectories can result in unpredictable lodgement sites and associated dental trauma; retained tooth fragments may mimic neoplasms on clinical exam, necessitating CT for diagnosis.^{2,3,7} Case 2 demonstrates that ethmoidal retention poses risk of orbital or intracranial extension; CT and endoscopic techniques enable safe removal with minimal morbidity.^{4,6,10}

CT remains the diagnostic gold standard; intraoperative imaging verifies complete retrieval. FESS is now preferred for sinus foreign bodies due to superior visualization and lower morbidity compared with open approaches. Adjunctive septoplasty may enhance access in anatomically restricted fields. Finally, prolonged mucosal contact with metallic debris carries the risk of lead absorption—clinicians should monitor lead levels when removal is incomplete.¹²

CONCLUSION

These two cases emphasize the rarity and complexity of retained ballistic foreign bodies in the upper aerodigestive tract. Timely CT imaging, multidisciplinary coordination, and individualized surgical strategies—whether open or endoscopic—are paramount to achieving favorable outcomes and preventing complications.

REFERENCES

1. Slusky E. A bullet in the tongue. *Oral Surg Oral Med Oral Pathol.* 1961;14(11):1394–1398.
2. Jiang ZH, Xv R, Xia L. Foreign body granuloma in the tongue differentiated from tongue cancer: a case report. *World J Clin Cases.* 2022;10(18):6247–6253.
3. Eliçora SŞ, Güven M. Delayed diagnosis of foreign body on the tongue: case report. *Braz J Otorhinolaryngology.* 2012;78(6):125.
4. Velasco GC, Araneta JMP, Añonuevo Cruz CEM. Bullet slug in the ethmoid sinus: a case report. *Philipp J Otolaryngol Head Neck Surg.* 2021;36(2):40–43.
5. Sharma RC. Gunshot injuries of maxillofacial region: current concepts in management. *Med J Armed Forces India.* 2005;61(3):238–241.
6. Kaufman Y, Buch K, Mahabadi N, Irish J. Facial gunshot wounds: a case series and review of management. *Craniofac Trauma Reconstr.* 2020;13(2):112–118.
7. Wężyk A, Paślawska U, Wróbel B, Gołda R, Niemczyk K. Metallic foreign body of the tongue: diagnostic and therapeutic approach. *Pol J Radiol.* 2016;81:325–328.
8. Viljoen G, Vermeulen MJ, Olorunju SAS. Orbito cranial gunshot injuries: experience from a South African trauma unit. *S Afr J Surg.* 2020;58(3):144–149.
9. Shanti RM, Egbert M, Sullivan SR. Management of penetrating injuries to the oral and maxillofacial region. *Oral Maxillofac Surg Clin North Am.* 2015;27(4):491–500.
10. Pandey N, Roychoudhury A, Trikha A, Bhutia O. Penetrating maxillofacial injuries due to projectile trauma: current concepts and clinical management. *J Oral Biol Craniofac Res.* 2015;5(3):193–198.
11. Lee D, Nash M, Turk J, HarEl G. Low velocity gunshot wounds to the paranasal sinuses. *Otolaryngology Head Neck Surg.* 1997;116(3):372–378.
12. Linden MA, Manton WI, Stewart RM, Thal ER, Feit H. Lead poisoning from retained bullets: pathogenesis, diagnosis, and management. *Ann Surg.* 1982;195(3):305–313. doi:10.1097/00000658-198203000-00010